



CONSISTENTLY POWERFUL RELIEF FOR YOUR KID'S ALLERGIES*

Children's ZYRTEC® Allergy Syrup*†



Use products only as directed.

Age	Dose
Adults and children 6 years and over	5 mL or 10 mL once daily depending upon severity of symptoms. Do not take more than 10 mL in 24 hours.
Children 2 to under 6 years	2.5 mL once daily. If needed, dose can be increased to a maximum of 5 mL once daily or 2.5 mL every 12 hours. Do not give more than 5 mL in 24 hours.
Children under 2 years	Ask a doctor.
Consumers with liver or kidney disease	Ask a doctor.
Adults 65 years and over	5 mL once daily. Do not take more than 5 mL in 24 hours.

Children's ZYRTEC® Chewables*



Use products only as directed.

Age	Dose: 2.5 MG CHEWABLES	Dose: 10 MG CHEWABLES
Children 2 to under 6 years	Chew and swallow 1 tablet (2.5 mg) once daily; if needed, dose can be increased to a maximum of 2 tablets (5 mg) once daily or 1 tablet (2.5 mg) every 12 hours. Do not give more than 2 tablets (5 mg) in 24 hours.	Ask a doctor.
Adults and children 6 years and over	Chew and swallow 2 tablets (5 mg) or 4 tablets (10 mg) once daily depending upon severity of symptoms; do not take more than 4 tablets (10 mg) in 24 hours.	Chew and swallow 1 tablet (10 mg) once daily; do not take more than 1 tablet (10 mg) in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
Children under 2 years	Ask a doctor.	Ask a doctor.
Consumers with liver or kidney disease	Ask a doctor.	Ask a doctor.
Adults 65 years and over	Chew and swallow 2 tablets (5 mg) once daily; do not take more than 2 tablets (5 mg) in 24 hours.	Ask a doctor.

Children's ZYRTEC® Dissolve Tabs*



Use product only as directed.

Age	Dose
Adults and children 6 years and over	One 10 mg tablet once daily; do not take more than one 10 mg tablet in 24 hours. A 5 mg product may be appropriate for less severe symptoms.
Children under 6 years	Ask a doctor.
Consumers with liver or kidney disease	Ask a doctor.
Adults 65 years and over	Ask a doctor.

For additional information on allergy management, talk to your pediatrician.

See other side for dosing information on Children's BENADRYL®

mL = milliliter.

*Relieves sneezing; runny nose; itchy, watery eyes; and itching of the nose or throat.

†Use only with dosing cup provided with product.



THE BRAND MOMS TURN TO FIRST*

*When they first treat their children's upper respiratory allergies.



AGE	DOSE
6-11 years	Take 5 mL to 10 mL every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
2-5 years	Do not use unless directed by a doctor.
Under 2 years	Do not use.


AGE	DOSE
12+ years	Take 10 mL every 4 hours. Do not take more than 6 doses in 24 hours.
6-11 years	Take 5 mL every 4 hours. Do not take more than 6 doses in 24 hours.
4-5 years	Do not use unless directed by a doctor.
Under 4 years	Do not use.

AGE	DOSE
12+ years	2 to 4 chewable tablets (25 mg to 50 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
6-11 years	1 to 2 chewable tablets (12.5 mg to 25 mg) every 4 to 6 hours or as directed by a doctor. Do not take more than 6 doses in 24 hours.
2-5 years	Do not use unless directed by a doctor.
Under 2 years	Do not use.



Children's BENADRYL®
Allergy Liquid†‡

Use products only as directed.



Children's BENADRYL®
Allergy Plus Congestion†‡

Use product only as directed.



Children's BENADRYL®
Chewables†

Use product only as directed.

See other side for dosing information
on Children's ZYRTEC®

For additional information on allergy management, talk to your pediatrician.

mL = milliliter.

†Relieves runny nose; sneezing; itchy, watery eyes; and itching of the nose or throat.

‡Use only with dosing cup provided with product.